

More congruences from Apéry-like formulae

ROBERTO TAURASO

Dipartimento di Matematica

Università di Roma “Tor Vergata”, Italy

tauraso@mat.uniroma2.it

<http://www.mat.uniroma2.it/~tauraso>

Abstract

We present some congruences modulo p^{6-r} for sums of the type $\sum_{k=0}^{(p-3)/2} x^k \binom{2k}{k} / (2k+1)^r$, for $r = 1, 2, 3$ where $p > 5$ is a prime. This is a preliminary draft and more refined versions will follow.

1 Introduction

In proving the irrationality $\zeta(3)$, Apéry [1] mentioned the identities

$$\sum_{k=1}^{\infty} \frac{1}{k^2} \binom{2k}{k}^{-1} = \frac{1}{3} \zeta(2), \quad \sum_{k=1}^{\infty} \frac{(-1)^k}{k^3} \binom{2k}{k}^{-1} = -\frac{2}{5} \zeta(3).$$

Later Koecher [4], and Leshchiner [5], found several analogous results for other $\zeta(r)$. In particular, in [5], the author gives several proofs of four identities. Two of them, namely (3b) and (4b), are connected with the series expansion of odd powers of the complex function $\arcsin(z)$ (see [2]): for $|z| < 2$ and for any odd integer $r > 0$ then

$$\arcsin(z/2)^r = \frac{r!}{2} \sum_{k=0}^{\infty} \frac{\binom{2k}{k}}{16^k} \cdot \frac{z^{2k+1}}{(2k+1)} \cdot \overline{H}_k(\{2\}^{\frac{r-1}{2}}).$$

where

$$\overline{H}_n(a_1, a_2, \dots, a_d) = \sum_{0 \leq k_1 < k_2 < \dots < k_r < n} \frac{1}{(2k_1+1)^{a_1} (2k_2+1)^{a_2} \dots (2k_r+1)^{a_r}},$$

with $(a_1, a_2, \dots, a_r) \in (\mathbb{N}^*)^r$.

By revisiting the combinatorial proof (due to D. Zagier) presented in Section 5 in [5], it is easy to obtain the finite versions of these identities: if r is a positive odd integer then

$$\begin{aligned} & \sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{16^k} \left(\sum_{j=0}^{\frac{r-1}{2}} \frac{(-1)^j \overline{H}_k(\{2\}^j)}{(2k+1)^{r-2j}} - \frac{(-1)^{\frac{r-1}{2}}}{4} \cdot \frac{\overline{H}_k(\{2\}^{\frac{r-1}{2}})}{(2k+1)} \right) \\ &= \sum_{k=0}^{n-1} \frac{(-1)^k}{(2k+1)^r} + \frac{(-1)^{\frac{r-1}{2}}}{4} \sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{16^k \binom{n+k}{2k+1}} \cdot \frac{(-1)^{n-k} \overline{H}_k(\{2\}^{\frac{r-1}{2}})}{(2k+1)}, \end{aligned} \quad (1)$$

and if r is a positive even integer then

$$\begin{aligned} \sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{(-16)^k} & \left(\sum_{j=0}^{\frac{r}{2}-1} \frac{(-1)^j \overline{H}_k(\{2\}^j)}{(2k+1)^{r-2j}} + \frac{(-1)^{\frac{r}{2}-1}}{4} \cdot \frac{\overline{H}_k(\{2\}^{\frac{r}{2}-1})}{(2k+1)^2} \right) \\ & = \sum_{k=0}^{n-1} \frac{1}{(2k+1)^r} + \frac{(-1)^{\frac{r}{2}-1}}{4} \sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{(-16)^k \binom{n+k}{2k+1}} \cdot \frac{\overline{H}_k(\{2\}^{\frac{r}{2}-1})}{(2k+1)^2}. \end{aligned} \quad (2)$$

After some preliminary results, starting from (1) for $r = 1$, and (2) for $r = 2$, that is

$$\frac{3}{4} \sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{16^k (2k+1)} = \sum_{k=0}^{n-1} \frac{(-1)^k}{(2k+1)} + \frac{(-1)^n}{4} \sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{(-16)^k \binom{n+k}{2k+1}} \cdot \frac{1}{2k+1}, \quad (3)$$

$$\frac{5}{4} \sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{(-16)^k (2k+1)^2} = \sum_{k=0}^{n-1} \frac{1}{(2k+1)^2} + \frac{1}{4} \sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{(-16)^k \binom{n+k}{2k+1}} \cdot \frac{1}{(2k+1)^2}, \quad (4)$$

we prove that for any prime $p > 5$

$$\begin{aligned} \sum_{k=0}^{\frac{p-3}{2}} \frac{\binom{2k}{k}}{16^k (2k+1)} & \equiv (-1)^{\frac{p-1}{2}} \left(\frac{H_{p-1}(1)}{12} + \frac{3}{160} p^4 B_{p-5} \right) \pmod{p^5}, \\ \sum_{k=0}^{\frac{p-3}{2}} \frac{\binom{2k}{k}}{(-16)^k (2k+1)^2} & \equiv \frac{H_{p-1}(1)}{5p} + \frac{7}{20} p^3 B_{p-5} \pmod{p^4}, \end{aligned}$$

where $H_n(r) = \sum_{k=1}^n \frac{1}{k^r}$ and B_n denotes the n -th Bernoulli number. In the last section we consider the case $r = 3$. These congruences confirm as conjectures in [11, Conjecture 5.1]. Notice that by taking the limit as n goes to infinity in (3) and (4) we obtain

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{\binom{2k}{k}}{16^k (2k+1)} & = \frac{4}{3} \sum_{k=0}^{\infty} \frac{(-1)^k}{(2k+1)} = \frac{4}{3} \arctan(1) = \frac{\pi}{3}, \\ \sum_{k=0}^{\infty} \frac{\binom{2k}{k}}{(-16)^k (2k+1)^2} & = \frac{4}{5} \sum_{k=0}^{\infty} \frac{1}{(2k+1)^2} = \frac{3}{5} \zeta(2) = \frac{\pi^2}{10}, \end{aligned}$$

which allow us to evaluate the analogy between the finite and the infinite sum. For more results of this flavour, involving Apéry-like formulae, see for example [8, 6, 12, 11]. In the recent preprint [9], the authors prove congruences modulo p^{4-r} for sums of the general form $\sum_{k=0}^{(p-3)/2} \binom{2k}{k} x^k / (2k+1)^r$ for $r = 1, 2$ in terms of the finite polylogarithms.

2 Results concerning multiple harmonic sums

We define the *multiple harmonic sum* as

$$H_n(a_1, a_2, \dots, a_r) = \sum_{0 < k_1 < k_2 < \dots < k_r \leq n} \frac{1}{k_1^{a_1} k_2^{a_2} \dots k_r^{a_r}},$$

where $n \geq r > 0$ and $(a_1, a_2, \dots, a_r) \in (\mathbb{N}^*)^r$. The values of many harmonic sums modulo a power of prime p are well known. Here there is a list of results that we will need later.

(i) ([10, Theorem 5.1]) for any prime $p > r + 2$ we have

$$H_{p-1}(r) \equiv \begin{cases} -\frac{r(r+1)}{2(r+2)} p^2 B_{p-r-2} \pmod{p^3} & \text{if } r \text{ is odd,} \\ \frac{r}{r+1} p B_{p-r-1} \pmod{p^2} & \text{if } r \text{ is even;} \end{cases}$$

(ii) ([13, Theorem 3.1]) for $r, s > 0$, and for any prime $p > r + s$, we have

$$H_{p-1}(r, s) \equiv \frac{(-1)^s}{r+s} \binom{r+s}{s} B_{p-r-s} \pmod{p};$$

(iii) ([13, Theorem 3.5]) for $r, s, t > 0$, and for any prime $p > r + s + t$ such that $r + s + t$ is odd, we have

$$H_{p-1}(r, s, t) \equiv \frac{1}{2(r+s+t)} \left((-1)^r \binom{r+s+t}{r} - (-1)^t \binom{r+s+t}{t} \right) B_{p-r-s-t} \pmod{p};$$

(iv) ([12, Theorem 2.1]) for any prime $p > 5$,

$$H_{p-1}(1) \equiv -\frac{1}{2} H_{p-1}(2) - \frac{1}{6} H_{p-1}(3) \pmod{p^5};$$

(v) ([8, Lemma 3]) for any prime $p > 5$,

$$H_{p-1}(1, 2) \equiv -3 \frac{H_{p-1}(1)}{p^2} + \frac{1}{2} p^2 B_{p-5} \pmod{p^3};$$

(vi) ([10, Theorem 5.2]) for any prime $p > r + 4$ we have

$$H_{\frac{p-1}{2}}(r) \equiv \begin{cases} -\frac{2^r-2}{r} B_{p-r} \pmod{p} & \text{if } r > 1 \text{ is odd,} \\ \frac{r(2^{r+1}-1)}{2(r+1)} p B_{p-r-1} \pmod{p^2} & \text{if } r \text{ is even.} \end{cases}$$

The rest of this section is dedicated to proving some congruences involving $H_{\frac{p-1}{2}}(r)$ and $H_{\frac{p-1}{2}}(r, s)$.

Lemma 2.1. *Let $r, a > 0$ then for any prime $p > r + 2$*

$$H_{p-1}(r) \equiv H_{\frac{p-1}{2}}(r) + (-1)^r \sum_{k=0}^a \binom{r-1+k}{k} H_{\frac{p-1}{2}}(r+k) p^k \pmod{p^{a+1}}, \quad (5)$$

$$\begin{aligned} H_{\frac{p-1}{2}}(r) &\equiv H_{\lfloor \frac{p}{4} \rfloor}(r) + (-2)^r \sum_{k=0}^a \binom{r-1+k}{k} H_{\frac{p-1}{2}}(r+k) p^k \\ &\quad - (-1)^r \sum_{k=0}^a \binom{r-1+k}{k} \frac{H_{\lfloor \frac{p}{4} \rfloor}(r+k)}{2^k} p^k \pmod{p^{a+1}}. \end{aligned} \quad (6)$$

Moreover, if $r, s > 0$ such that $r + s$ is odd then for any prime $p > r + s$

$$H_{\frac{p-1}{2}}(r, s) \equiv \frac{B_{p-r-s}}{2(r+s)} \left((-1)^s \binom{r+s}{s} + 2^{r+s} - 2 \right) \pmod{p}. \quad (7)$$

Proof. The congruence (5) follows from the identity

$$H_{p-1}(r) = H_{\frac{p-1}{2}}(r) + \sum_{k=1}^{\frac{p-1}{2}} \frac{1}{k^r(1-p/k)^r}.$$

Moreover the identity

$$H_{\frac{p-1}{2}}(r) = H_{\lfloor \frac{p}{4} \rfloor}(r) + 2^r \left(\sum_{k=1}^{(p-1)/2} \frac{1}{(p-k)^r} - \sum_{k=1}^{\lfloor p/4 \rfloor} \frac{1}{(p-2k)^r} \right)$$

yields (6). Now we show (7).

$$\begin{aligned} H_{p-1}(r, s) &= \sum_{1 \leq i < j \leq \frac{p-1}{2}} \frac{1}{i^r j^s} + \sum_{1 \leq i, j \leq \frac{p-1}{2}} \frac{1}{i^r (p-j)^s} + \sum_{1 \leq j < i \leq \frac{p-1}{2}} \frac{1}{(p-i)^r (p-j)^s} \\ &\equiv H_{\frac{p-1}{2}}(r, s) + (-1)^s H_{\frac{p-1}{2}}(r) H_{\frac{p-1}{2}}(s) + (-1)^{r+s} H_{\frac{p-1}{2}}(s, r) \\ &\equiv H_{\frac{p-1}{2}}(r, s) - H_{\frac{p-1}{2}}(s, r) \pmod{p}. \end{aligned}$$

By the stuffle product property

$$H_{\frac{p-1}{2}}(r, s) + H_{\frac{p-1}{2}}(s, r) + H_{\frac{p-1}{2}}(r+s) = H_{\frac{p-1}{2}}(r) H_{\frac{p-1}{2}}(s) \equiv 0 \pmod{p}.$$

Therefore

$$2H_{\frac{p-1}{2}}(r, s) \equiv H_{p-1}(r, s) - H_{\frac{p-1}{2}}(r+s) \pmod{p}$$

and by applying (ii) and (vi) we obtain (6). $\square$

Theorem 2.2. For any prime $p > 2$

$$H_{\frac{p-1}{2}}(2) + \frac{7}{6} p H_{\frac{p-1}{2}}(3) + \frac{5}{8} p^2 H_{\frac{p-1}{2}}(4) \equiv 0 \pmod{p^4}. \quad (8)$$

Proof. Let $m = \varphi(p^4) = p^3(p-1)$ and let $B_n(x)$ the n -th Bernoulli polynomial. For $r = 2, 3, 4$, Faulhaber's formula implies

$$\begin{aligned} \sum_{k=1}^{(p-1)/2} k^{m-r} &= \frac{B_{m-r+1}\left(\frac{p-1}{2}\right) - B_{m-r+1}}{m-r+1} \\ &= \frac{B_{m-r+1}\left(\frac{p-1}{2}\right) - B_{m-r+1}\left(\frac{1}{2}\right)}{m-r+1} + \frac{B_{m-r+1}\left(\frac{1}{2}\right) - B_{m-r+1}}{m-r+1} \\ &= \sum_{k=r}^m \binom{m-r}{k-r} \frac{B_{m-k}\left(\frac{1}{2}\right)}{k-r+1} \cdot \left(\frac{p}{2}\right)^{k-r+1} + \frac{B_{m-r+1}\left(\frac{1}{2}\right) - B_{m-r+1}}{m-r+1}. \end{aligned}$$

By Euler's theorem, $H_{\frac{p-1}{2}}(r) \equiv \sum_{k=1}^{(p-1)/2} k^{m-r} \pmod{p^4}$. Since m is even, $B_{m-k} = 0$ when $m-k > 1$ and k is odd. Moreover, pB_{m-k} is p -integral and

$$B_{m-k}\left(\frac{1}{2}\right) = (2^{1-m+k} - 1)B_{m-k} \equiv (2^{1+k} - 1)B_{m-k} \pmod{p^4}.$$

Hence

$$\begin{aligned} \sum_{r=2}^4 \alpha_r p^{r-2} H_{\frac{p-1}{2}}(r) &\equiv \frac{p}{2} \left(\alpha_2 - \frac{6}{7} \alpha_3 \right) B_{m-2} \left(\frac{1}{2} \right) \\ &\quad + \frac{p^3}{8} (\alpha_2 - 3\alpha_3 + 4\alpha_4) B_{m-4} \left(\frac{1}{2} \right) \pmod{p^4}. \end{aligned}$$

The right-hand side vanishes if we let $\alpha_2 = 1$, $\alpha_3 = 7/6$, and $\alpha_4 = 5/8$. $\square$

Corollary 2.3. *For any prime $p > 5$*

$$H_{p-1}(2) \equiv -2 \frac{H_{p-1}(1)}{p} + \frac{2}{5} p^3 B_{p-5} \pmod{p^4}, \quad (9)$$

$$H_{\frac{p-1}{2}}(2) \equiv -7 \frac{H_{p-1}(1)}{p} + \frac{17}{10} p^3 B_{p-5} \pmod{p^4}, \quad (10)$$

$$H_{\frac{p-1}{2}}(3) \equiv 6 \frac{H_{p-1}(1)}{p^2} - \frac{81}{10} p^2 B_{p-5} \pmod{p^3}, \quad (11)$$

$$H_{\frac{p-1}{2}}(1, 2) + p H_{\frac{p-1}{2}}(1, 3) \equiv -\frac{9}{2} \frac{H_{p-1}(1)}{p^2} - \frac{49}{20} p^2 B_{p-5} \pmod{p^3}. \quad (12)$$

Proof. The congruence (9) follows from (iv) and (i). By (8) and (vi) we have

$$H_{\frac{p-1}{2}}(2) + \frac{7}{6} p H_{\frac{p-1}{2}}(3) + \frac{31}{4} B_{p-5} p^3 \equiv 0 \pmod{p^4}.$$

From (5) and (vi), we deduce

$$H_{p-1}(2) = 2H_{\frac{p-1}{2}}(2) + 2pH_{\frac{p-1}{2}}(3) + \frac{66}{5} B_{p-5} p^3 \equiv 0 \pmod{p^4}.$$

By solving these two congruences with respect to $H_{\frac{p-1}{2}}(2)$ and $H_{\frac{p-1}{2}}(3)$, and by using (9), we get the result. Now we show (12). We consider the identity

$$H_{p-1}(1, 2) = H_n(1, 2) + H_n(1) \sum_{j=1}^n \frac{1}{(p-j)^2} + \sum_{1 \leq j < i \leq n} \frac{1}{(p-i)(p-j)^2}$$

and by expanding the sums like in Lemma 2.1, we get

$$\begin{aligned} H_{p-1}(1, 2) &\equiv H_n(1, 2) + H_n(1)(H_n(2) + 2p H_n(3) + 3p^2 H_n(4)) - H_n(2, 1) \\ &\quad - p(2H_n(3, 1) + H_n(2, 2)) - p^2(3H_n(4, 1) + 2H_n(3, 2) + H_n(2, 3)) \pmod{p^3}. \end{aligned}$$

By applying the stuffle product for $H_n(1)H_n(2)$ and $H_n(1)H_n(3)$, and (i), (ii) and (vi), we obtain

$$H_n(1, 2) + p H_n(3, 1) \equiv \frac{1}{2} H_{p-1}(1, 2) - \frac{1}{2} H_n(3) - \frac{27}{4} p^2 B_{p-5} \pmod{p^3}.$$

Thus the proof of (12) is complete as soon as we apply (v) and (11). $\square$

Theorem 2.4. *For any prime $p > 5$*

$$\begin{aligned} 2(-1)^{\frac{p-1}{2}} \sum_{k=0}^{\frac{p-3}{2}} \frac{(-1)^k}{2k+1} &\equiv \overline{H}_{\frac{p-1}{2}}(1) - p \overline{H}_{\frac{p-1}{2}}(2) - p^2 \overline{H}_{\frac{p-1}{2}}(2, 1) \\ &\quad + p^3 \overline{H}_{\frac{p-1}{2}}(2, 2) + p^4 \overline{H}_{\frac{p-1}{2}}(2, 2, 1) \pmod{p^5}, \end{aligned} \quad (13)$$

$$\equiv -\frac{1}{2} H_{\frac{p-1}{2}}(1) + \frac{11}{16} H_{p-1}(1) - \frac{57}{1280} p^4 B_{p-5} \pmod{p^5}. \quad (14)$$

Proof. The following identities hold

$$\sum_{k=0}^n \frac{(-16)^k \binom{n+k}{2k}}{(2k+1) \binom{2k}{k}} = 2(-1)^n \sum_{k=0}^{n-1} \frac{(-1)^k}{2k+1} + \frac{1}{2n+1} \quad \text{and} \quad \sum_{k=0}^n \frac{(-16)^k \binom{n+k}{2k}}{(2k+1)^2 \binom{2k}{k}} = \frac{1}{(2n+1)^2}.$$

Let $n = (p-1)/2$. Notice that for $k = 0, \dots, n$

$$\frac{(-16)^k \binom{n+k}{2k}}{\binom{2k}{k}} = \prod_{j=0}^{k-1} \left(1 - \frac{p^2}{(2j+1)^2} \right) = \sum_{j=0}^{k-1} (-1)^j p^{2j} \overline{H}_k(\{2\}^j). \quad (15)$$

Thus the two identities yield

$$\begin{aligned} \overline{H}_n(1) - p^2 \overline{H}_n(2, 1) + p^4 \overline{H}_n(2, 2, 1) &\equiv 2(-1)^n \sum_{k=0}^{n-1} \frac{(-1)^k}{2k+1} + \frac{1 - (-1)^n 4^{p-1} \binom{2n}{n}^{-1}}{p} \pmod{p^6}, \\ \overline{H}_n(2) - p^2 \overline{H}_n(2, 2) + p^4 \overline{H}_n(2, 2, 2) &\equiv \frac{1 - (-1)^n 4^{p-1} \binom{2n}{n}^{-1}}{p^2} \pmod{p^6}, \end{aligned}$$

and, by subtracting p times the second congruence from the first one, we get (13).

As regards (14), let $m = \lfloor p/4 \rfloor$, then

$$\begin{aligned} (-1)^n \sum_{k=0}^{n-1} \frac{(-1)^k}{2k+1} &= \sum_{k=1}^n \frac{(-1)^k}{p-2k} = \sum_{k=1}^m \frac{1}{p-4k} - \sum_{k=1}^m \frac{1}{p-(4k-2)} \\ &= 2 \sum_{k=1}^m \frac{1}{p-4k} - \sum_{k=1}^n \frac{1}{p-2k} \equiv -\frac{1}{2} \sum_{k=0}^4 \frac{H_m(k+1)}{4^k} p^k + \frac{1}{2} \sum_{k=0}^4 \frac{H_n(k+1)}{2^k} p^k \pmod{p^5}. \end{aligned}$$

By taking a suitable linear combination of the equations (6) for $r = 1, 2, 4$ we find that

$$\sum_{k=0}^4 \frac{H_m(k+1)}{4^k} p^k \equiv \frac{3}{2} H_n(1) + \frac{13}{16} p H_n(2) + \frac{1}{2} p^2 H_n(3) + \frac{143}{512} p^3 H_n(4) + \frac{1}{8} p^4 H_n(5) \pmod{p^5}.$$

Hence

$$(-1)^n \sum_{k=0}^{n-1} \frac{(-1)^k}{2k+1} \equiv -\frac{1}{4} H_n(1) - \frac{5}{32} p H_n(2) - \frac{1}{8} p^2 H_n(3) - \frac{79}{1024} p^3 H_n(4) - \frac{1}{32} p^4 H_n(5) \pmod{p^5},$$

and (14) follows by applying (10), (11), and (vi). $\square$

Note that by using (13), (14) and (iii), it can be shown that for any prime $p > 5$,

$$H_{\frac{p-1}{2}}(2, 1, 2) \equiv 2 H_{\frac{p-1}{2}}(1, 2, 2) \equiv 10 H_{\frac{p-1}{2}}(2, 2, 1) \equiv -\frac{15}{4} B_{p-5} \pmod{p}.$$

Moreover, another consequence of the previous theorem is a generalization of Morley's congruence [3]: for any prime $p > 5$,

$$\frac{(-1)^{\frac{p-1}{2}}}{4^{p-1}} \binom{p-1}{\frac{p-1}{2}} \equiv 1 - \frac{1}{4} p H_{p-1}(1) - \frac{1}{80} p^5 B_{p-5} \pmod{p^6}.$$

3 Proof of the main result

We are finally ready to prove the congruences announced in the introduction.

Theorem 3.1. *For any prime $p > 5$ we have*

$$\sum_{k=0}^{\frac{p-3}{2}} \frac{\binom{2k}{k}}{16^k(2k+1)} \equiv (-1)^{\frac{p-1}{2}} \left(\frac{H_{p-1}(1)}{12} + \frac{3}{160} p^4 B_{p-5} \right) \pmod{p^5}, \quad (16)$$

$$\sum_{k=0}^{\frac{p-3}{2}} \frac{\binom{2k}{k}}{(-16)^k(2k+1)^2} \equiv \frac{H_{p-1}(1)}{5p} + \frac{7}{20} p^3 B_{p-5} \pmod{p^4}. \quad (17)$$

Proof. From (15), it follows that

$$\begin{aligned} \frac{\binom{2k}{k}}{(-16)^k \binom{n+k}{2k+1}} &= \frac{(2k+1)\binom{2k}{k}}{(-16)^k(n-k)\binom{n+k}{2k}} \equiv -\frac{2}{\left(1 - \frac{p}{2k+1}\right) (1 - p^2 \overline{H}_k(2) + p^4 \overline{H}_k(4) - p^6 \overline{H}_k(6))} \\ &\equiv -2 \left(1 + \frac{p}{2k+1} + \left(\frac{1}{(2k+1)^2} + \overline{H}_k(2) \right) p^2 + \left(\frac{1}{(2k+1)^3} + \frac{\overline{H}_k(2)}{2k+1} \right) p^3 \right. \\ &\quad \left. \left(\frac{1}{(2k+1)^4} + \frac{\overline{H}_k(2)}{(2k+1)^2} + \overline{H}_k(4) + \overline{H}_k(2, 2) \right) p^4 \right) \pmod{p^5}. \end{aligned} \quad (18)$$

Therefore, by (3), (18) and (13), we get

$$\begin{aligned} \frac{3(-1)^n}{2} \sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{16^k(2k+1)} &\equiv -2p \overline{H}_n(2) - p^2 (\overline{H}_n(3) + 2\overline{H}_n(2, 1)) - p^3 \overline{H}_n(4) \\ &\quad - p^4 (\overline{H}_n(5) + \overline{H}_n(2, 3) + \overline{H}_n(4, 1)) \pmod{p^5}. \end{aligned} \quad (19)$$

Now we replace the terms $\overline{H}_n$ with the corresponding expressions involving H_{p-1} and H_n :

$$\begin{aligned} \overline{H}_n(r) &= H_{p-1}(r) - \frac{H_n(r)}{2^r}, \\ \overline{H}_n(r, s) &= \sum_{0 < j < i \leq n} \frac{1}{(2(n-i)+1)^r (2(n-j)+1)^s} = \frac{1}{(-2)^{r+s}} \sum_{0 < j < i \leq n} \frac{1}{j^s i^r \left(1 - \frac{p}{2i}\right)^r \left(1 - \frac{p}{2j}\right)^s} \\ &\equiv \frac{1}{(-2)^{r+s}} \left(H_n(s, r) + \frac{p}{2} (r H_n(s, r+1) + s H_n(s+1, r)) \right. \\ &\quad \left. + \frac{p^2}{4} \left(\binom{r+1}{2} H_n(s, r+2) + r s H_n(s+1, r+1) + \binom{s+1}{2} H_n(s+2, r) \right) \right) \pmod{p^3}. \end{aligned}$$

So the right-hand side of (19) becomes

$$\begin{aligned} &-2p H_{p-1}(2) - p^2 H_{p-1}(3) - p^3 H_{p-1}(4) - p^4 H_{p-1}(5) \\ &+ \frac{p}{2} H_n(2) + \frac{p^2}{8} (H_n(3) + 2H_n(1, 2)) + \frac{p^3}{16} (H_n(4) + 4H_n(1, 3) + 2H_n(2, 2)) \\ &- \frac{p^4}{32} (H_n(5) + 4H_n(2, 3) + 3H_n(3, 2) + 7H_n(1, 4)) \pmod{p^5} \end{aligned}$$

Since

$$H_n(2, 2) = \frac{1}{2} (H_n(2)^2 - H_n(4)) \equiv -\frac{31}{5} p B_{p-5} \pmod{p^2},$$

by (i), (ii), and (vi), the above expression simplifies to

$$-2pH_{p-1}(2) + \frac{1}{2}pH_n(2) + \frac{1}{8}p^2H_n(3) + \frac{1}{4}p^2(H_n(1,2) + pH_n(1,3)) + \frac{513}{320}p^4B_{p-5} \pmod{p^5}.$$

Finally we apply (9), (10), (11) and (12)

$$\frac{H_{p-1}(1)}{8} + \frac{9}{320}p^4B_{p-5} \pmod{p^5}$$

which concludes our proof of (16).

As regards (17), by (4) and (18), we have

$$\begin{aligned} 5 \sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{(2k+1)^2(-16)^k} &\equiv 4\overline{H}_n(2) - 2(\overline{H}_n(2) + p\overline{H}_n(3) + p^2(\overline{H}_n(4) + \overline{H}_n(2,2)) \\ &\quad + p^3(\overline{H}_n(5) + \overline{H}_n(2,3))) \pmod{p^4}. \end{aligned} \quad (20)$$

As before, after replacing the terms $\overline{H}_n$, the right-hand side of (20) becomes

$$\begin{aligned} 2H_{p-1}(2) - 2pH_{p-1}(3) - 2p^2H_{p-1}(4) - 2p^3H_{p-1}(5) \\ - \frac{1}{2}H_n(2) + \frac{p}{4}H_n(3) + \frac{p^2}{8}(H_n(4) - H_n(2,2)) + \frac{p^3}{16}(H_n(5) - 2H_n(2,3) - H_n(3,2)) \pmod{p^4}. \end{aligned}$$

By (i), (ii), and (vi), it simplifies to

$$2H_{p-1}(2) - \frac{1}{2}H_n(2) + \frac{1}{4}pH_n(3) + \frac{9}{4}p^3B_{p-5} \pmod{p^4}$$

Finally we apply (9), (10) and (11) and we get

$$\frac{H_{p-1}(1)}{p} + \frac{7}{4}p^3B_{p-5} \pmod{p^4}.$$

□

4 The case $r = 3$

From (1) for $r = 3$, we have

$$\sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{16^k} \left(\frac{1}{(2k+1)^3} - \frac{3}{4} \frac{\overline{H}_k(2)}{(2k+1)} \right) = \sum_{k=0}^{n-1} \frac{(-1)^k}{(2k+1)^3} - \frac{(-1)^n}{4} \sum_{k=0}^{n-1} \frac{\binom{2k}{k} \overline{H}_k(2)}{(-16)^k \binom{n+k}{2k+1}} \cdot \frac{1}{2k+1}. \quad (21)$$

Let $n = (p-1)/2$ and let $m = \lfloor p/4 \rfloor$, then, by taking a suitable linear combination of the equations (6) for $r = 3, 5$ we find that

$$\begin{aligned} (-1)^n \sum_{k=0}^{n-1} \frac{(-1)^k}{(2k+1)^3} &= 2 \sum_{k=1}^m \frac{1}{(p-4k)^3} - \sum_{k=1}^n \frac{1}{(p-2k)^3} \\ &\equiv -\frac{1}{32} \left(H_m(3) + \frac{3H_m(3)}{4}p + \frac{3H_m(4)}{8}p^2 \right) \\ &\quad + \frac{1}{8} \left(H_n(3) + \frac{3H_n(3)}{2}p + \frac{3H_n(4)}{2}p^2 \right) \\ &= -\frac{1}{64}H_n(3) - \frac{3}{16}pH_n(4) - \frac{189}{512}p^2H_n(5) \\ &\equiv -\frac{3}{32} \frac{H_n(1)}{p^2} + \frac{21}{1280}p^2B_{p-5} \pmod{p^3}. \end{aligned}$$

Moreover, acting as in Theorem 3.1, we obtain that

$$\begin{aligned} \sum_{k=0}^{n-1} \frac{\binom{2k}{k} \overline{H}_k(2)}{(-16)^k \binom{n+k}{2k+1}} \cdot \frac{1}{2k+1} &\equiv 2\overline{H}_n(2, 1) + 2p\overline{H}_n(2, 2) \\ &\quad + 2p^2(\overline{H}_n(2, 3) + \overline{H}_n(4, 1) + 2\overline{H}_n(2, 2, 1)) \\ &\equiv -\frac{9}{8} \frac{H_{p-1}(1)}{p^2} + \frac{9}{320} p^2 B_{p-5} \pmod{p^3}. \end{aligned}$$

Hence

$$(-1)^n \sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{16^k} \left(\frac{1}{(2k+1)^3} - \frac{3}{4} \frac{\overline{H}_k(2)}{(2k+1)} \right) \equiv \frac{3}{16} \frac{H_{p-1}(1)}{p^2} + \frac{3}{320} p^2 B_{p-5} \pmod{p^3}. \quad (22)$$

On the other hand, since 3 does not divide $2n+1$, by (15), we have

$$0 = \sum_{k=0}^{n-1} \binom{n+k}{2k+1} \frac{(-1)^k}{2k+1} \equiv \sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{16^k (2k+1)} (1 - p^2 \overline{H}_k(2) + p^4 \overline{H}_k(2, 2)) \pmod{p^5}.$$

Since (the proof will be given in the next version)

$$\sum_{k=0}^{n-1} \frac{\binom{2k}{k} \overline{H}_k(2, 2)}{16^k (2k+1)} \equiv (-1)^n \frac{5}{864} B_{p-5} \pmod{p},$$

by (16), it follows that

$$\sum_{k=0}^{n-1} \frac{\binom{2k}{k} \overline{H}_k(2)}{16^k (2k+1)} \equiv (-1)^n \left(\frac{H_{p-1}(1)}{12p^2} + \frac{53}{2160} p^2 B_{p-5} \right) \pmod{p^3}. \quad (23)$$

Finally, by (22), we get

$$\sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{16^k (2k+1)^3} \equiv (-1)^n \left(\frac{H_{p-1}(1)}{4p^2} + \frac{1}{36} p^2 B_{p-5} \right) \pmod{p^3}. \quad (24)$$

Note that the values of the corresponding infinite series are known (see [7]):

$$\sum_{k=0}^{n-1} \frac{\binom{2k}{k}}{16^k (2k+1)^3} = \frac{7\pi^3}{216} \quad \text{and} \quad \sum_{k=0}^{\infty} \frac{\binom{2k}{k} \overline{H}_k(2)}{16^k (2k+1)} = \frac{\pi^3}{648}.$$

References

- [1] R. Apéry, *Irrationalité de $\zeta(2)$ et $\zeta(3)$* . *Journées arithmétiques de Luminy*, Astérisque **61** (1979), 11–13.
- [2] J. M. Borwein and M. Chamberland, *Integer powers of arcsin*, Int. J. Math. Math. Sciences (2007), Art. ID 19381.
- [3] L. Carlitz, *A theorem of Glaisher*, Canadian J. Math. **5** (1953), 306–316.
- [4] M. Koecher, *Letter (German)*, Math. Intelligencer, **2** (1979/1980), no. 2, 62–64.

- [5] D. H. Leshchiner, *Some new identities for $\zeta(k)$* , J. Number Theory **13** (1981), 355–362.
- [6] S. Mattarei, R. Tauraso, *Congruences for central binomial sums and finite polylogarithms*, preprint, arXiv:1012.1308v6 (2011).
- [7] Kh. Hessami Pilehrood, T. Hessami Pilehrood, *Series acceleration formulas for beta values*, Discrete Math. Theor. Comput. Sci. **12** (2010), 223–236.
- [8] Kh. Hessami Pilehrood, T. Hessami Pilehrood, *Congruences arising from Apéry-type series for zeta values*, preprint, arXiv:1108.1893v2 (2011).
- [9] Kh. Hessami Pilehrood, T. Hessami Pilehrood, *Congruences concerning Jacobi polynomials*, preprint, arXiv:1110.5308v1 (2011).
- [10] Z. H. Sun, *Congruences concerning Bernoulli numbers and Bernoulli polynomials*, Discrete Appl. Math. **105** (2000), 193–223.
- [11] Z. W. Sun, *On congruences related to central binomial coefficients*, J. Number Theory **131** (2011), 2219–2238.
- [12] R. Tauraso, *More congruences for central binomial coefficients*, J. Number Theory **130** (2010), 2639–2649.
- [13] J. Zhao, *Wolstenholme type theorem for multiple harmonic sums*, Int. J. Number Theory **4** (2008), 73–106.